

Ventilatory Mechanics in Early vs Late Intubation in a Cohort of COVID-19 Patients With Acute Respiratory Distress Syndrome

A Single Center's Experience

To the Editor:

Coronavirus disease 2019 (COVID-19) is associated with a range of presentations, from milder symptoms to severe hypoxic respiratory failure, often meeting criteria for ARDS. Patients admitted to an ICU are likely to require mechanical ventilation (up to 85% in US cohorts), which is associated with increased morbidity and mortality.^{1,2}

Methods

This retrospective study includes adult inpatients requiring invasive mechanical ventilation secondary to COVID-19 at Temple University Hospital between February and May 2020. Positive infection status was confirmed by polymerase chain reaction nasopharyngeal swab. The study was derived from the institutional review board (IRB)-approved Temple University Registry for COVID-19 (TUIRB Protocol Number: 26854). Subsequently, a separate IRB approval was granted for chart review to extract ventilator settings (TUIRB protocol number: 27051).

Data were collected from the electronic medical record and managed using REDCap electronic data capture tools from the Temple University Hospital COVID-19 Registry.³

Results

Seventy-five patients with nasopharyngeal swab-confirmed COVID-19 required invasive mechanical ventilation at Temple University Hospital during the study period. Average age was 65 years, and median BMI was 31.8. Fifty-eight percent of the patients were male, and 63% were African American.

Median time to intubation was 1.27 days from presentation. Patients were separated into an early intubation (≤ 1.27 days) or late intubation (> 1.27 days) group for analysis.

Patients in the late intubation group had a lower F_{iO_2} requirement on admission (55% vs 69%; $P = .109$), but a

The near universal approach to early mechanical ventilation at the onset of the COVID-19 pandemic was driven by early data from China describing rapid deterioration with severe hypoxia, fears of patient self-induced lung injury and infection control measures because of concern about aerosolization in non-intubated patients. There remains a paucity of literature describing respiratory mechanics, ventilatory parameters, and outcomes in relation to early and late intubation in COVID-19 patients.

We report the ventilatory parameters and lung mechanics of consecutive early and late intubated and ventilated patients with COVID-19 ARDS by descriptive analysis at a single urban academic center in Philadelphia, Pennsylvania.

All patients met Berlin criteria for ARDS. Patients required invasive mechanical ventilation for acute hypoxic respiratory failure based on the P_{aO_2}/F_{iO_2} ratio, or by clinical decision. Lung protective ventilation strategies and adjunct therapies were employed per general guidelines and physician discretion. Daily recorded ventilator parameters were analyzed. Individual patient lung compliance, driving pressures, and ventilatory ratios (VR) were calculated.⁴

Descriptive statistics, either in mean with SD or median with interquartile range (IQR) and percentages, were used to present clinical data. Significance testing between groups was done with Student t test or Wilcoxon rank sum with continuous data or χ^2 with categorical data.

worse P_{aO_2}/F_{iO_2} ratio (median, 160 vs 205; $P = .46$), higher PEEP (11.29 vs 9.30; $P = .027$), plateau (26.41 vs 22.50; $P = .027$), and peak pressures (32.21 vs 28.62; $P = .044$) at time of intubation in comparison with the early intubation group.

Lower static compliance (34.88 vs 40.68; $P = .311$) and higher VR (1.90 vs 1.57; $P = .078$) was noted in the late intubation group on day 0, although these values were not statistically significant. Static compliance increased by day 6 of intubation in the late group, whereas it decreased in those intubated early (39.80 vs 31.66; $P = .129$). The late intubation group did have a significantly longer length of stay in the ICU (median, 12.31

TABLE 1] Patient Characteristics, Respiratory Support and Parameters, Ventilatory Parameters, and Outcomes in Early vs Late Intubation for COVID-19 ARDS

| Variable | NMiss | Time to Intubation | | | χ^2 or <i>t</i> Test P Value |
|--|-------|---------------------------|---------------------------|---------------------------------|--------------------------------------|
| | | Total (n = 75) | Early (<1.27) (n = 37) | Late (\geq 1.27) (n = 38) | |
| Age, No. | 0 | 75 | 37 | 38 | .575 |
| Mean (SD) | | 64.97 (14.27) | 65.92 (14.79) | 64.05 (13.87) | |
| BMI, No. | 2 | 73 | 35 | 38 | .002 |
| Median (IQR) | | 31.80 (25.83-39.48) | 28.63 (22.61-35.35) | 34.16 (29.52-41.02) | |
| Sex, No. (%) | 0 | | | | .921 |
| Female | | 32 (42.67) | 16 (50.00) | 16 (50.00) | |
| Male | | 43 (57.33) | 21 (48.84) | 22 (51.16) | |
| Race, No. (%) | 0 | | | | .291 |
| African American | | 47 (62.67) | 22 (46.81) | 25 (53.19) | |
| Caucasian | | 8 (10.67) | 3 (37.50) | 5 (62.50) | |
| Hispanic | | 17 (22.67) | 9 (52.94) | 8 (47.06) | |
| Other | | 3 (4.00) | 3 (100.00) | 0 (0.00) | |
| Time to intubation, No. | 0 | 75 | 37 | 38 | <.0001 |
| Mean (SD) | | 2.86 (4.47) | 0.15 (0.30) | 5.51 (5.03) | |
| Initial F _{IO₂} at time of admission, No. | 21 | 54 | 25 | 29 | .109 |
| Mean (SD) | | 0.62 (0.32) | 0.69 (0.31) | 0.55 (0.33) | |
| Pao ₂ /F _{IO₂} at intubation, No. | 4 | 71 | 34 | 37 | .460 |
| Median (IQR) | | 162.00 (106.00-316.00) | 205.50 (106.00-378.00) | 160.00 (99.00-268.00) | |
| Positive end expiratory pressure, No. | 0 | 75 | 37 | 38 | .027 |
| Mean (SD) | | 10.31 (3.93) | 9.30 (3.75) | 11.29 (3.89) | |
| Plateau pressure, No. | 16 | 59 | 30 | 29 | .027 |
| Mean (SD) | | 24.42 (6.74) | 22.50 (4.76) | 26.41 (7.91) | |
| Peak pressure, No. | 0 | 75 | 37 | 38 | .044 |
| Mean (SD) | 16 | 30.44 (7.73) | 28.62 (6.75) | 32.21 (8.29) | |
| Driving pressure, No. | 16 | 59 | 30 | 29 | .154 |
| Mean (SD) | | 14.36 (6.12) | 13.23 (4.77) | 15.53 (7.16) | |
| Static compliance day 0, No. | 16 | 59 | 30 | 29 | .311 |
| Mean (SD) | | 37.83 (21.95) | 40.68 (27.23) | 34.88 (14.59) | |
| Static compliance day 6, No. | 41 | 34 | 14 | 20 | .129 |
| Mean (SD) | | 36.45 (16.87) | 31.66 (10.16) | 39.80 (19.87) | |
| Ventilatory ratio at intubation, No. | 6 | 69 | 32 | 37 | .078 |
| Mean (SD) | | 1.75 (0.78) | 1.57 (0.63) | 1.90 (0.86) | |
| Duration of ventilation, No. | 33 | 42 | 22 | 20 | .102 |
| Mean (SD) | | 7.98 (8.77) | 5.86 (8.40) | 10.30 (8.78) | |
| ICU length of stay, days, No. | 0 | 75 | 37 | 38 | .001 |
| Median (IQR) | | 9.25 (5.42-16.25) | 7.38 (3.88-10.21) | 12.31 (7.75-19.96) | |
| Length of stay, days, No. | 0 | 75 | 37 | 38 | .037 |
| Median (IQR) | | 13.00 (4.00-19.00) | 10.00 (1.00-15.00) | 15.50 (8.00-22.00) | |
| Living status, No. (%) | 0 | | | | .563 |
| Deceased | | 37 (49.33) | 17 (45.95) | 20 (54.05) | |
| Living | | 38 (50.67) | 20 (52.63) | 18 (47.37) | |

vs 7.38 days; $P = .001$) and duration of mechanical ventilation (10.30 vs 5.86; $P = .102$) (Table 1).

As of data censoring on June 20, 2020, 49% of all mechanically ventilated patients had died. The median age for nonsurvivors was higher than those for survivors (70 vs 59; $P = .0006$). Average time to intubation was 3.88 days in nonsurvivors and 1.87 in survivors ($P = .053$). Nonsurvivors had higher initial FiO_2 requirement (70% vs 50%; $P = .139$), lower $\text{PaO}_2/\text{FiO}_2$ ratio (median, 146 vs 261; $P = .010$), lower static compliance (32.14 vs 34.62; $P = .962$), and higher ventilatory ratios (1.85 vs 1.64; $P = .276$).

Discussion

Our study found late intubation (>1.27 days; median, day 4) was associated with longer ICU length of stay and longer duration of mechanical ventilation than early intubation (≤ 1.27 days; median, day 0). We found that nonsurvivors had a longer time to intubation than survivors in our cohort.

Patients intubated later had higher driving pressures, lower static compliance, and higher ventilatory ratios. By day 6, static compliance improved in the late intubation group, whereas it declined in the early intubation group. This may be partially explained by disease improvement over time. Additionally, not all patients were included in the static compliance measures by day 6, because several patients had been extubated or expired. Low static compliance was seen in both groups of patients, albeit at varying times during the mechanical ventilation course. We did not find distinct ARDS phenotypes as previously suggested,⁵ in line with results from other cohort studies, suggesting that most patients have low compliance.⁶⁻⁹

This study has numerous limitations, including its retrospective nature. Only patients who were polymerase chain reaction positive were included. The decision to intubate was based on clinician preference; thus, time to intubation varied. The late intubation group had a significantly longer need for mechanical ventilation and time in the ICU. Although respiratory mechanics seemingly improved in this group, our study does not account for other causes and co-morbidities that may have contributed to prolonged mechanical ventilation. Furthermore, causes of mortality were not fully analyzed. Larger cohort studies are needed to detect a difference in mortality between early and late intubation.

In our cohort, all ventilated COVID-19 patients had low compliance and increased ventilatory ratios. Patients intubated later during their hospitalization appear to have

worse compliance or VR with potentially higher mortality. Whether this is progression of disease or the presence of patient self-induced lung injury remains unclear. Further studies will need to be performed to determine whether onset of symptoms, time to hospitalization, timing of intubation, and pharmacotherapies are variables that can alter a patient's clinical course.

Alok Nath Pandya, MD

Navjot Ariyana Kaur, MD

Daniel Sacher, DO

Oisin O'Corragain, MD

Daniel Salerno, MD

Parag Desai, MD

Sameep Sehgal, MD

Matthew Gordon, MD

Rohit Gupta, MD

Nathaniel Marchetti, DO

Huaqing Zhao, PhD

Nicole Patlakh, BSc

Gerard J. Criner, MD

Temple University COVID-19 Research Group*

Philadelphia, PA

AFFILIATIONS: From the Department of Thoracic Medicine and Surgery, Lewis Katz School of Medicine at Temple University.

FINANCIAL/NONFINANCIAL DISCLOSURES: The authors have reported to CHEST the following: G. J. C. reports grants from Fisher and Paykel, during the conduct of the study; grants and personal fees from Galaxo Smith Kline, grants and personal fees from Boehringer Ingelheim, grants and personal fees from Chiesi, grants and personal fees from Mereo, personal fees from Verona, grants and personal fees from Astra Zeneca, grants and personal fees from Pulmonx, grants and personal fees from Pneumrx, personal fees from BTG, grants and personal fees from Olympus, grants and personal fees from Broncus, personal fees from EOLO, personal fees from NGM, grants and personal fees from Lungpacer, grants from Alung, grants and personal fees from Nuaira, grants and personal fees from ResMed, grants and personal fees from Respiromics, grants from Fisher Paykel, grants and personal fees from Patara, grants from Galapagos outside the submitted work. None declared (A. P., N. A. K., D. S., O. O. C., D. S., P. D., S. S., M. G., R. G., N. M., N. P.).¹

*Collaborators from the Temple University COVID-19 Research Group are listed in the e-Appendix.

CORRESPONDENCE TO: Navjot Ariyana Kaur, MD, Department of Thoracic Medicine and Surgery, Lewis Katz School of Medicine at Temple University, Philadelphia, PA; e-mail: navjot.ariyana@gmail.com

Copyright © 2020 American College of Chest Physicians. Published by Elsevier Inc. All rights reserved.

DOI: <https://doi.org/10.1016/j.chest.2020.08.2084>

Acknowledgments

Additional information: The e-Appendix can be found in the Supplemental Materials section of the online article.

References

1. Bhatraju PK, Ghassemieh BJ, Nichols M, et al. Covid-19 in critically ill patients in the Seattle region: case series. *N Engl J Med*. 2020;382(21):2012-2022.

276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330

Q9

Q1 Q2
Q4 Q3

Q5

Q8

Q10

Q6

- 331 2. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting
332 characteristics, comorbidities, and outcomes among 5700 patients
333 hospitalized with COVID-19 in the New York City area. *JAMA*.
334 2020;323(20):2052-2059. 386
- 335 3. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG.
336 Research electronic data capture (REDCap): A metadata-driven
337 methodology and workflow process for providing translational
338 research informatics support. *J Biomed Inform*. 2009;42(2):377-381. 387
- 339 4. Sinha P, Fauvel NJ, Singh P, Soni N. Analysis of ventilatory ratio as a
340 novel method to monitor ventilatory adequacy at the bedside. *Crit
341 Care*. 2013;17(1):R34. 388
- 342 5. Marini JJ, Gattinoni L. Management of COVID-19 respiratory
343 distress [Published online ahead of print April 24, 2020]. *JAMA*. 2020.
344 <https://doi.org/10.1001/jama.2020.6825>. 389
- 345 6. Bos LD, Paulus F, Vlaar APJ, Beenen LFM, Schultz MJ.
346 Subphenotyping ARDS in COVID-19 patients: consequences for
347 ventilator management. *Ann Am Thorac Soc*. 2020;17(9):1161-1163. 390
- 348 7. Schenck EJ, Hoffman K, Goyal P, et al. Respiratory mechanics and gas
349 exchange in COVID-19 associated respiratory failure. *Ann Am
350 Thorac Soc*. 2020;17(9):1158-1161. 391
- 351 8. Ziehr DR, Alladina J, Petri CR, et al. Respiratory pathophysiology of
352 mechanically ventilated patients with COVID-19: a cohort study. *Am
353 J Respir Crit Care Med*. 2020;201(12):1560-1564. 392
- 354 9. Gattinoni L, Coppola S, Cressoni M, Busana M, Rossi S,
355 Chiumello D. COVID-19 does not lead to a “typical” acute
356 respiratory distress syndrome. *Am J Respir Crit Care Med*.
357 2020;201(10):1299-1300. 393
- 358 394
- 359 395
- 360 396
- 361 397
- 362 398
- 363 399
- 364 400
- 365 401
- 366 402
- 367 403
- 368 404
- 369 405
- 370 406
- 371 407
- 372 408
- 373 409
- 374 410
- 375 411
- 376 412
- 377 413
- 378 414
- 379 415
- 380 416
- 381 417
- 382 418
- 383 419
- 384 420
- 385 421